

# Management of subependymal giant cell tumors in tuberous sclerosis complex: the neurosurgeon's perspective

Moncef Berhouma

Lyon, France

**Background:** Tuberous sclerosis complex (TSC), an autosomal dominant genetic disorder, can lead to the development of hamartomas in various organs, including the heart, lungs, kidneys, skin and brain. The management of subependymal giant cell tumors (SGCTs) is still controversial, and peri- and/or intraventricular neoplasms may lead to life-threatening hydrocephalus. In the last years, many progresses have been made in research into the tumorigenesis and behaviors of SGCTs. This review aims to clarify the specific role of neurosurgeons in the multidisciplinary management of SGCTs in children with TSC.

**Data sources:** Based on the recent scientific literature and personal experience, we reviewed the up-to-date data and discussed the trends in the management of SGCTs in children with TSC. The data were collected after a bibliography made using PubMed/Medline with these terms: subependymal, subependymal giant cell astrocytoma, subependymal giant cell tumor, and tuberous sclerosis complex.

**Results:** SGCTs are shown to be generated from a glioneuronal lineage, but their filiation with subependymal nodules (SENs) is still under debate. While SENs may develop anywhere in the ventricular walls, SGCTs arise almost exclusively around the Monro foramina. In children with TSC, precise clinical and/or imaging criteria are mandatory to differentiate SENs that are always asymptomatic and riskless from SGCTs that have the potential to grow and therefore to obstruct cerebrospinal fluid pathways leading to hydrocephalus.

**Conclusions:** An earlier diagnosis of SGCT in neurologically asymptomatic children with TSC may

allow a precocious surgical removal of the tumor before the installation of increased intracranial pressure signs, an attitude that is being progressively adopted to lessen the morbimortality rate.

*World J Pediatr* 2010;6(2):103-110

**Key words:** cerebral ventricle; hydrocephalus; intraventricular tumor; microsurgery; subependymal giant cell astrocytoma; tuberous sclerosis

## Introduction

Tuberous sclerosis complex (TSC) is an autosomal dominant genetic disorder, with a low and variable penetrance, characterized by the formation of multiple tumors in different organs. The disease is caused by mutations in either the *TSC1* gene (Chromosome 9) or the *TSC2* gene (Chromosome 16) which encodes hamartin and tuberin respectively. These proteins form an intracellular complex involved in the regulation of the cellular growth and energetic pathways, through the inhibition of the mammalian target of rapamycin (mTOR) within the akt-mTOR-S6 kinase cell growth trail. TSC, also known historically as Bourneville's disease (1880), has been initially described by von Recklinghausen in 1862, while Vogt (1908) reported the typical and characteristic clinical triad of the disease associating facial angiofibromatosis, epilepsy and mental retardation.<sup>[1,2]</sup>

TSC is one of the more common single gene disorders, with an incidence of one in 5800 to 6000 live births, a prevalence of one in 30 000 and a frequency of one per 150 000 live births.<sup>[3,4]</sup> There is neither ethnic linking nor differences according to gender.<sup>[5]</sup> In spite of all the progresses carried out in the comprehension of the pathogenesis of the disease, TSC remains a disabling situation for the affected child and his entourage in terms of social and educational integration. The disease usually leads to epilepsy, learning difficulties and behavioral anomalies. It is broadly admitted that about 50% of children with TSC have an intelligence quotient within a

**Author Affiliations:** Department of Neurosurgery B (Unit 501), Pierre Wertheimer Hospital, Lyon, France; Claude Bernard Lyon 1 University, France (Berhouma M)

**Corresponding Author:** Moncef Berhouma, MD, Associate Professor, Department of Neurosurgery B, Pierre Wertheimer Hospital, 59 Boulevard Pinel 69394 Lyon Cedex 03, France (Tel: 0033684284713; Email: berhouma.moncef@yahoo.fr)

doi:10.1007/s12519-010-0025-2

©Children's Hospital, Zhejiang University School of Medicine, China and Springer-Verlag Berlin Heidelberg 2010. All rights reserved.

normal range while the other half will usually suffer from severe cognitive impairment.<sup>[4]</sup> Among the different sites of tumor development, the brain remains undoubtedly the most problematic regarding to therapeutic management and screening, as this location is directly related to life expectancy. This location leads to more than 50% of deaths among children with TSC.<sup>[6]</sup> Since many years, several interrogations are pending especially concerning subependymal giant cell tumors (SGCTs) in children with TSC: cytological origin and pathological classification, probable filiation with subependymal nodules (SENs), diagnosis criteria and screening, and neurosurgical indications.<sup>[7-11]</sup> Indeed, while symptomatic SGCTs in children with TSC do not require any debate concerning their surgical management, asymptomatic ones remain more problematic to handle even if a recent more "interventionist" tendency is emanating from the neurosurgical literature.<sup>[1,3,9,12,13]</sup> Thus, after a brief reminder of the main clinical features of TSC and a compilation of the current data concerning the pathogenesis of TSC and the management of SGCTs, the author tries to clarify the modest role of the neurosurgeon in the multidisciplinary management of SGCTs in children with TSC.

### Clinical background

An accurate diagnosis of TSC is based on precise criteria resulting from consensus conferences.<sup>[14,15]</sup> Roughly, the diagnosis is retained when at least two hamartomas are found in two different organs in an individual. Because of the variable penetrance of the disease, children with TSC may display a wide range of intelligence levels. Generally it is admitted that 50% of TSC children have a normal intelligence quotient while the other half suffers from severe learning difficulties, leading to social and educational marginalization.<sup>[16]</sup> Besides, three main types of lesions might be encountered during the evolution of the disease: cortical tubers, SENs and SGCTs. Firstly, tubers are developmental cortical abnormalities of the brain, present in more than 88% of children with TSC.<sup>[3]</sup> Tubers lead to the loss of the classical six-layered cyto-architecture of the cerebral cortex. These lesions are better detected on MRI. Classically, they appear as hypointense lesions in T1 sequence and hyperintense in T2 sequence.<sup>[3,17,18]</sup> These frequent lesions are thought to be responsible for more than 75% of epileptic disorders in patients with TSC.<sup>[3,14]</sup> During the first year of life, infantile spasms are common and respond habitually to GABA inhibitors especially vigabatrin.<sup>[19]</sup> Later in life, protean seizures including atonic, focal or atypical fits are generally refractory to medical treatments. These can benefit nowadays from epilepsy surgery since it is technically possible

to identify clearly epileptogenic tubers with the use of functional imaging and cortical mapping techniques.<sup>[20,21]</sup> The second more frequent lesions in children with TSC are SENs. These are small hamartomas developed on the walls of the lateral ventricles. There is no evidence that SENs can cause any neurological symptoms. Only SENs located in the region of the Monro foramina may have the potentiality to grow and to transform into SGCTs. The last but not least encephalic lesion is SGCT, also known improperly as subependymal giant cell astrocytoma, affecting an average of 10% of children with TSC. The filiation between SENs and SGCTs is still a great matter of debate. The main enigma remains to understand why SGCTs develop almost exclusively near the Monro foramina. In addition to these neurological manifestations, TSC children may display other potentially lethal systemic locations (heart, lungs, kidneys).<sup>[22-25]</sup>

### Genetic acquirements

Despite TSC is an autosomal dominant genetic disease, about 70% of cases result from new mutations (unaffected parents). Otherwise, the gonadal mosaicism phenomenon applies to TSC, i.e., an apparently unaffected parent has a population of affected cells confined to his gonads, so that unaffected parents may have more than one affected child.<sup>[26]</sup> More than 15 years ago, linkage studies have individualized the 2 genes responsible for TSC, namely *TSC1* (chromosome 9) and *TSC2* (chromosome 16).<sup>[27-29]</sup> A mutation in either of the 2 genes might induce all the manifestations of the disease. Meanwhile, two facts have been noted. First, *TSC2* phenotype related disease seems to be more severe.<sup>[30]</sup> Second, the frequency of mutations appears to be higher in the *TSC2* gene.<sup>[31]</sup> In spite of the improvements made in the genetic field, only 80% of TSC cases are being detected during the prenatal screening because of the occurrence of somatic mosaicism. Hence, an affected child has 2 populations of cells, i.e., one population expresses a genetic mutation (*TSC1* or *TSC2*) while the other pool of cells does not.<sup>[32]</sup> At a molecular level, *TSC1* product is hamartin, a 140 kDa protein expressed in several adult tissues and playing a key role in the regulation of cell adhesion. Tuberin (*TSC2* product) is a 200 kDa protein involved in the GTPase activating protein activity.<sup>[33]</sup> Hamartin and tuberin act together in the akt-mTOR-S6 kinase cell growth pathway, interfering directly in the regulation of cell growth, differentiation and proliferation.<sup>[34]</sup> Recent insights in the comprehension of the disease set the course to new trails in the therapeutic possibilities. Consequently, the use of drugs that may counterbalance the deficit of hamartin or tuberin seems tempting. It is the case with

rapamycin, a well-known immunosuppressor that is also thought to be an inhibitor of the mTOR and therefore would theoretically inhibit the S6kinase activity in children with TSC. Rapamycin has been successfully used to shrink SGCTs in patients with TSC but the lack of sufficient data concerning the requisite duration of treatment and the existence of side effects (aphtous ulcerations, interstitial pneumonitis) restrict its common use in children with TSC at the present time.<sup>[35,36]</sup> Further studies to determine the duration of treatment and the risk of tumor recurrence after discontinuation of rapamycin are certainly needed.

### Tumorigenesis and behavior of SGCTs

For many decades, SGCT has been improperly classified among astrocytomas even in the last WHO classification of central nervous system tumors. This terminology definitively merits revision. Recent pathological studies including molecular biology, electron microscopy and immunochemistry categorically rectified this erroneous nomenclature, confirming the glioneuronal lineage of SGCTs.<sup>[12]</sup> This explains why the term SGCT is actually being preferred to the widely used "subependymal giant cell astrocytoma". Cytological features are highly characteristic and of an important interest in differentiating SGCTs from other intraventricular neoplasms mainly ependymomas and gemistocytic astrocytomas: cell clustering, hypercellularity and fibrillarity.<sup>[12]</sup> Immunochemistry contributes also to grounding the diagnosis as SGCTs are glial fibrillary acidic protein, neuron specific enolase, synaptophysin, and neurofilament positive. In the meantime, histological and immunohistochemical data do not provide any useful information on the behavior of SGCTs. There is a complete lack of correlation between the tumoral course and the pathological features.<sup>[12,37-39]</sup> Thus, an increased mitotic index, pleiomorphism, endothelial proliferation and necrosis seem to have no prognostic value in SGCTs.<sup>[3,12]</sup>

It has been noted that MIB-1 labeling indices are low in these neoplasms even in recurrences.<sup>[12,40]</sup> It has also been established that there is no correlation between kinetic index (Ki-67) and SGCTs growth.<sup>[3,12]</sup> As the tumorigenesis is concerned, the development of hamartomas in TSC fits the two-hit model presented by Knudson:<sup>[41]</sup> the first hit corresponds to a congenital lesion of either *TSC1* or *TSC2*, and the second hit is a loss of heterozygosity (LOH) of this gene. This model applies perfectly to most of the neoplasms in TSC, but only 30%-60% of SGCTs and cardiac rhabdomyomas show effectively LOH. This latter is particularly rare in SGCTs.<sup>[42,43]</sup> Therefore, it has been hypothesized that SGCTs might share a common feature mimicking

LOH, corresponding probably to an inactivation of TSC by a phosphorylation process or to a direct activation of mTOR through two mighty protein kinases: protein kinase B (AKT) and extra-cellular signal-regulated kinase (ERK). While the first one is not regularly detected, the second seems to be hyperactive in all SGCTs.<sup>[7]</sup> Jozwiak et al<sup>[7]</sup> postulated recently that ERK activation might be a molecular trigger for the development of SGCTs. Despite all the recent data based on molecular biology concerning the tumorigenesis in children with TSC, we still do not understand why the different hamartomas associated with TSC do not appear at the same age of the evolution of the disease and have flagrant discrepancies in growth profile. For example, renal angiomyolipomas grow rapidly and usually appear in adolescents and young adults. Cardiac rhabdomyomas emerge during fetal life and generally regress at birth, while SGCTs slowly but rarely become symptomatic in adults.<sup>[22,44]</sup>

### Filiations SEN-SGCT

SGCTs are estimated to occur in about 6.1% to 18.5% of children with TSC.<sup>[3,6,8,38]</sup> The fact that these tumors arise from pre-existing SENs is still a great matter of debate. This hypothesis lies not only on growth evidence of SENs shown by serial imaging studies, but also on the pathological similarity between the two lesions.<sup>[9,17,45]</sup> The LOH and/or a protein phosphorylation process has been put forth as trigger factors in the transformation phase of SENs into SGCTs.<sup>[46-48]</sup> For this filiation SEN-SGCT, we consider why SGCTs arise almost exclusively around the foramen of Monro when SENs may appear anywhere in the lateral ventricle walls? Up to now, no valid explanation has been proposed, while 88% to 95% of patients with TSC present SENs,<sup>[9,45]</sup> and about one child with TSC out of ten will develop SGCT(s). Exceptionally, SGCTs have been described in other ventricular compartments than the foramen of Monro, such as the lateral ventricle wall<sup>[2]</sup> or the temporal horn.<sup>[49]</sup> Elsewhere, extremely rare cases of neonatal SGCTs have contributed to our incomprehension of tumorigenesis and call in question again the supposed filiation SEN-SGCT.<sup>[50]</sup> Distinguishing SENs from SGCTs is essential, since it is mostly admitted that SENs do not grow and never induce clinical signs, while SGCTs have the potential to enlarge and become symptomatic. Therefore, the therapist needs to identify which SEN should be diagnosed as a tumor in order to take effect early. Clinical and radiological criteria to differentiate these two entities are still in debate.<sup>[1,3,10,51]</sup> SENs are totally asymptomatic lesions, whereas the existence of raised intracranial pressure (ICP) signs, visual disturbances, focal neurological signs, or a worsening of epilepsy

suggest the presence of SGCT(s) in children with TSC.

It is also admitted that SGCTs are exceptionally revealed in adults.<sup>[6,12]</sup> Thus no new SGCTs are diagnosed after the age of 21 even if a known SGCT may continue to grow in adulthood.<sup>[10]</sup> Nowadays, with the multiplication of MRI centers, a majority of children with TSC are diagnosed earlier before the apparition of the above-mentioned neurological signs and symptoms. In such cases, radiological criteria are important in distinguishing SENs from SGCTs. Thus, a subependymal expansive lesion measuring more than 5 mm of diameter, developed immediately around the foramen of Monro, containing fine calcifications and displaying frank contrast enhancement associated with hydrocephalus, is more likely to be a SGCT.<sup>[1]</sup> The author emphasizes the growth evidence on serial MRI as important distinctive criteria rather than a fixed and arbitrary diameter of the lesion (Table).

## Neurosurgical management of SGCTs

### Timing of surgery

In the past only symptomatic children with TSC were operated on (tumor removal and/or ventriculoperitoneal shunt), and there is actually a general belief that a more precocious surgical intervention provides a lesser morbidity or mortality than when increased ICP or hydrocephalus is already installed, i.e., surgery in

asymptomatic children.<sup>[1,10,12,52]</sup> SGCTs represent about 1.4% of all pediatric brain tumors.<sup>[3,37]</sup> Clinical series showed a mild male predominance, with a mean age at surgery of 11 years.<sup>[3,10,13,17]</sup> Habitually, these neoplasms evolve on several weeks or months. They are revealed by increased ICP symptoms, epilepsy recrudescence and behavioral modifications. Exceptionally, a severe onset may disclose the tumor through an acute hydrocephalus, an intraventricular hemorrhage or even a sudden death.<sup>[53,54]</sup> Actually, a rationale trend advocates an earlier tumoral removal in asymptomatic TSC children, an attitude from which we easily understand the importance of the distinctive criteria between SENs and SGCTs. This surgical procedure proposed to neurologically asymptomatic TSC children amounts to discuss the hazards of the so-called preventive surgery. This situation reminds the similar dilemma of an incidental colloid cyst of the third ventricle discovered by an imaging done for head injury. Nevertheless, several established facts strengthen this proceeding:

First, serial imaging studies have already proven that SGCTs have the potential to grow and therefore to become symptomatic. Because of their preferential proximity with the foramen of Monro, SGCTs may obstruct cerebrospinal fluid (CSF) pathways or cause intraventricular hemorrhage, and in both cases they can lead to death. In contrast, SENs do not possess this growth potentiality. In Cuccia's series, all SENs (about 70 nodules), except those near the foramen of Monro, remained unchanged or even decreased in size.<sup>[3]</sup>

Second, it has been stated that the growth of SGCTs may worsen epileptic seizures in terms of frequency and intensity, probably through the associated hydrocephalus and/or the direct irritation of the interventricular septum.<sup>[1]</sup>

Third, large SGCTs induce a deformation of the foramen of Monro, leading to difficulties in operative hemostasis and problems of tumoral dissection from surrounding structures such as the fornix, the head of caudate nucleus, the ependyma and its veins, and the interventricular septum. For this reason, it is obvious that operating on smaller lesions might avoid these surgical drawbacks.

Fourth, it is proven that major complications occur in operated symptomatic children with TSC when increased ICP is already installed.<sup>[1,10,12]</sup>

For all these reasons, it seems rational to recommend early tumoral resection as soon as a subependymal expansive lesion shows the distinctive criteria of SGCTs in asymptomatic screened children with TSC, primarily when there is evidence of growth on successive MRI.

### Management of associated hydrocephalus

Hydrocephalus is present in the majority of children at

**Table.** Clinical and radiological distinctive criteria between subependymal nodules (SENs) and subependymal giant cell tumors (SGCTs)


| Criteria                    | SENs  | SGCTs  |
|-----------------------------|---|--|
| Clinical signs and symptoms | Asymptomatic  | - Raised intracranial pressure<br>- Epilepsy worsening (intensity and frequency of seizures)<br>- Neurological focal signs<br>- Visual disturbances<br>- Accentuation of cognitive impairment and behavioral changes |
| Disclosure age              | Asymptomatic  | Exceptionally revealed after 21 years <sup>[10]</sup>  |
| CT-scan                     | Hyperdense<br>Frequent pat calcifications                                   | - Isodense or slightly hyperdense<br>- Rare thin calcifications  |
| MRI                         | Variable signal depending on relaxation time<br>Slight contrast enhancement | - Hypointense in T1<br>- Hypointense in T2<br>- Marked contrast enhancement  |
| Growth on serial imaging    | Absent  | 3 to 4 mm per year <sup>[3]</sup>  |
| Diameter                    | Inferior to 5 mm  | Superior to 5 mm <sup>[1]</sup>  |
| Cerebral ventricles         | Normal  | Frequent hydrocephalus   |
| Predilection site           | Anywhere on the lateral ventricle walls                                     | Foramen of Monro   |

the moment of the diagnosis of SGCT. Twelve of the 15 patients of Cuccia's series and 7 of the 11 patients of Goh's one displayed hydrocephalus.<sup>[3,10]</sup> This is the main factor contributing to the increase of ICP rather than the tumoral volume itself. Nevertheless, increased ICP may be present even in the absence of ventriculomegaly in children with TSC, because of the abnormal architecture of the subependymal region, thus modifying the cerebral compliance.<sup>[10]</sup> The treatment of hydrocephalus lies on the surgical resection of the tumor that blocks the foramen of Monro. There is no place actually for the placement of a ventriculoperitoneal shunt before the tumor's removal, except in the rare cases of acute life-threatening hydrocephalus. Elsewhere, it appears that an abnormal high level of CSF proteins in non-removed SGCTs leads to the frequent obstruction of CSF diversion devices in shunted children. This CSF proteins level may normalize after resection of the tumor.<sup>[10,13]</sup> This last fact corroborates the uselessness of ventriculoperitoneal shunt in hydrocephalic children


with TSC before the resection of SGCTs. Finally, it is important to keep in mind that even after a complete removal of SGCTs, hydrocephalus can evolve on itself and even lead to postoperative deaths in 10% to 20% of removed SGCTs.<sup>[1,37]</sup> Following surgical resection of SGCTs, a rigorous postoperative clinical and radiological surveillance is mandatory to detect an eventual hydrocephalus in time (Fig. 1).

### Surgical technique and outcome

As SGCTs are benign lesions, complete removal remains the treatment of choice. This removal is synonymous with a nearly cure. These lesions develop almost always in the surroundings of the foramen of Monro, generally bulging within the lateral ventricle, sometimes in the third ventricle, nay in both (Fig. 2).<sup>[10,49,55]</sup> Surgical approach depends upon tumor extension and the presence of an associated hydrocephalus, but above all the surgeon's experience. Transcortical transventricular and transcallosal interhemispheric




**Fig. 1.** A: Axial T1-weighted image with gadolinium showing a bulky SGCT around the right Monro foramina, extending into the homolateral frontal horn with a cystic extension and a markedly contrast enhanced solid part. The tumor leads to hydrocephalus. Note also the presence of many calcified SENs on the walls of both lateral ventricles; B: Serial postoperative CT-scans of the same child (A) without contrast. Left: Immediate postoperative scan demonstrating a complete removal of the tumor performed through a right-sided transfrontal transventricular route, with some blood clots within the occipital horns. Middle and right: First postoperative day after a rapid neurological deterioration (Glasgow coma score 7/15), concomitant with a critical acute hydrocephalus requiring an urgent ventriculoperitoneal shunt, resulting in an immediate resolution of the ventriculomegaly and a complete neurological recovery.




**Fig. 2.** A: Coronal T1-weighted image without gadolinium: Isointense intraventricular mass developed in the right Monro foramina and extending in the right lateral ventricle as well as in the third ventricle, with a subsequent asymmetric hydrocephalus; B: Coronal T1-weighted image with contrast showing an intense and homogeneous contrast enhancement of the SGCT; C: Sagittal T1-weighted image with gadolinium: typical aspect of SGCT developed on horseback between the lateral ventricle and the third ventricle through a blocked right Monro foramina.


routes remain the most used approaches to the foramen of Monro, while endoscopic procedures are taking an increasing place among the surgical armamentarium of ventricular surgery, even in cases of small ventricles (stereotaxy).<sup>[1]</sup> Incomplete removal is generally the peculiarity of bulky SGCTs (Fig. 3) that lead to the deformation of local anatomy and a bad dissection plane (Fig. 4). The surgical outcome fluctuates between good and excellent in the majority of children operated on early. In Goh's series of 11 patients with a mean age of 11 years at surgery, an excellent surgical outcome was reported in 5 of the 8 preoperatively screened patients, while a poor outcome was noted for the 3 non-screened patients.<sup>[10]</sup> On the long term, complications occurred in all the patients who were older than 11 years at the time of surgery whereas this outcome was excellent in all children younger than 11 years. Goh's study suggests that young patients operated on have a better long-term outcome than older ones.<sup>[10]</sup> Other studies confirmed the benefits of an early surgical resection of SGCTs, i.e., when surgery is undertaken in young patients especially when tumor's diameter is less than 3 cm.<sup>[3,56]</sup> The complications after surgical removal of SGCTs rejoin those of any tumor surgery within the cerebral ventricles and around the foramen of Monro and its adjacent anatomical structures, namely the head of the caudate nucleus, the fornix and the ependymal veins (Fig. 5): transient hemiparesis (10%-12%),<sup>[3,49]</sup> permanent motor deficit (6%-12.5%),<sup>[3,10,37]</sup> reintervention for hemorrhage or compressive subdural collection (13%-20%).<sup>[1,3,57]</sup> An acute postoperative fatal hydrocephalus is not a rare event as it may occur in 10% to 20% of cases, generally secondary to infection or hemorrhage.<sup>[1,37,57]</sup> A postoperative cognitive impairment may also be observed, but problematical to assess as it occurs in




**Fig. 3.** A: Axial CT-scan without contrast demonstrating a left expansive intraventricular isodense and homogeneous mass obstructing the left Monro foramina associated with an important asymmetric hydrocephalus; B: Axial CT-scan with contrast showing an important but heterogeneous enhancement of the SGCT.



**Fig. 4.** A: Axial-T1 weighted image without gadolinium: SGCT blocking the left Monro foramina and leading to a unilateral ventriculomegaly; B: Axial-T1 weighted image with contrast: homogeneous enhancement of the mass with a heterogeneous bad-defined extension toward the head of the left caudate nucleus, predicting potential difficulties of surgical dissection.



**Fig. 5.** A: Operative photograph of a transcallosal interhemispheric approach showing clearly the corpus callosum (cc) before its section between the 2 pericallosal arteries (A); B: After a longitudinal callosotomy, the pursuit of the homolateral choroid plexus allows the surgeon to reach the right Monro foramina (MF), anteriorly edged by the fornix (Fx), laterally by the head of the caudate nucleus and medially by the interventricular septum.



**Fig. 6.** Suggested algorithm for the management of subependymal giant cell tumors (SGCTs).

young children with already preoperative cognitive anomalies and is not always imputable to a surgical damage to the fornix.<sup>[1]</sup> As previously stated, the surgical goal in the management of SGCTs is the complete and safe removal whenever possible, which means almost cure as no recurrences were noted at the end of 6.2 years of follow-up.<sup>[1]</sup> If the complete resection is not feasible because of excessive tumoral bleeding or the absence of a dissection plane, a close clinical and radiological follow-up is necessary with the following two situations (Fig. 6). Firstly, the residual tumor remains stable over years<sup>[12]</sup> but will require all the same a prolonged and close follow-up since late recurrences, sometimes fatal, have been reported.<sup>[3,58]</sup> Nevertheless, there is a general belief that SGCTs stop growing in adulthood even if it is not the rule.<sup>[55,58]</sup> Secondly, the residual tumor progresses on serial follow-up imaging. In this eventuality, a second surgery through the same approach or a different one may be proposed.<sup>[1,3,12,37,56]</sup> Another possibility relies on the use of the medical treatment (rapamycin) as this latter showed preliminary hopeful ascertainments resulting in tumoral shrinkage, but its side-effects and the duration of necessary treatment need to be evaluated in further studies.<sup>[35,36]</sup> During the coming years, medical treatment will certainly take more place in the management of children with TSC in proportion as the pathogenesis at the molecular level will be better understood.<sup>[59,60]</sup>

## Acknowledgement

The author thanks Prof. Hafedh Jemel for his constant scientific support.

**Funding:** None.

**Ethical approval:** Not needed.

**Competing interest:** None.

**Contributors:** Berhouma M is the single author of this paper.

## References

- Ribaupierre S, Dorfmueller G, Bulteau C, Fohlen M, Pinard JM, Chiron C, et al. Subependymal giant-cell astrocytomas in paediatric tuberous sclerosis disease: When should we operate? *Neurosurgery* 2007;60:83-90.
- Griffiths PD, Martland TR. Tuberous sclerosis complex: the role of neuroradiology. *Neuropediatrics* 1997;28:244-252.
- Cuccia V, Zuccaro G, Sosa F, Monges J, Lubienieky F, Taratuto A. Subependymal giant cell tumor in children with tuberous sclerosis. *Childs Nerv Syst* 2003;19:232-243.
- Osborne JP, Fryer A, Webb D. Epidemiology of tuberous sclerosis. *Ann N Y Acad Sci* 1991;615:125-127.
- Joinson C, O'Callaghan FJ, Osborne JP, Martyn C, Harris T, Bolton PF. Learning disability and epilepsy in an epidemiological sample of individuals with tuberous sclerosis complex. *Psychol Med* 2003;33:335-344.
- Webb DW, Fryer AE, Osborne JP. Morbidity associated with tuberous sclerosis: a population study. *Dev Med Child Neurol* 1996;38:146-155.
- Jozwiak J, Jozwiak S, Wlodarski P. Possible mechanisms of disease development in tuberous sclerosis. *Lancet Oncol* 2008;9:73-79.
- Roszkowski M, Drabik K, Barszcz S, Jozwiak S. Surgical treatment of intraventricular tumors associated with tuberous sclerosis. *Childs Nerv Syst* 1995;11:335-339.
- Nabbout R, Santos M, Rolland Y, Delalande O, Dulac O, Chiron C. Early diagnosis of subependymal giant cell tumor in children with tuberous sclerosis. *J Neurol Neurosurg Psychiatry* 1999;66:370-375.
- Goh S, Butler W, Thiele EA. Subependymal giant cell tumours in tuberous sclerosis complex. *Neurology* 2004;63:1457-1461.
- Narayanan V. Tuberous sclerosis complex: genetics to pathogenesis. *Pediatr Neurol* 2003;29:404-409.
- Kim SK, Wang KC, Cho BK, Jung HW, Lee YJ, Chung YS, et al. Biological behaviour and tumorigenesis of subependymal giant cell astrocytomas. *J Neurooncol* 2001;52:217-225.
- Di Rocco C, Iannelli A, Marchese E. On the treatment of subependymal giant cell astrocytomas and associated hydrocephalus in tuberous sclerosis. *Pediatr Neurosurg* 1995;23:115-121.
- Gomez MR. Criteria for diagnosis. In: Gomez MR, eds. *Tuberous sclerosis*. New York: Raven Press, 1998: 9-20.
- Hyman MH, Whittemore VH. National Institutes of Health consensus conference: tuberous sclerosis complex. *Arch Neurol* 2000;57:662-665.
- O'Callaghan FJ, Harris T, Joinson C, Bolton P, Noakes M, Presdee D, et al. The relation of infantile spasms, tubers, and intelligence in tuberous sclerosis complex. *Arch Dis Child* 2004;89:530-533.
- Braffman BH, Bilaniuk LT, Naidich TP, Altman NR, Post MJ, Quencer RM, et al. MR imaging of tuberous sclerosis: pathogenesis of this phakomatosis, use of gadopentetate dimeglumine, and literature review. *Radiology* 1992;183:227-238.
- Menor F, Martí-Bonmatí L, Mulas F, Poyatos C, Cortina H. Neuroimaging in tuberous sclerosis: a clinicoradiological evaluation in pediatric patients. *Pediatr Radiol* 1992;22:485-489.
- Hancock E, Osborne JP, Milner P. The treatment of West syndrome: a Cochrane review of the literature to December 2000. *Brain Dev* 2001;23:624-634.
- Romanelli P, Verdecchia M, Rodas R, Seri S, Curatolo P. Epilepsy surgery for tuberous sclerosis. *Pediatr Neurol* 2004;31:239-247.
- Madhavan D, Schaffer S, Yankovsky A, Arzimanoglou A, Renaldo F, Zaroff CM, et al. Surgical outcome in tuberous sclerosis complex: a multicenter survey. *Epilepsia* 2007;48:1625-1628.
- Jozwiak S, Kotulska K, Kasprzyk-obara J, Domańska-Pakieła D, Tomyn-Drabik M, Roberts P, et al. Clinical and genotype studies of cardiac tumors in 154 patients with tuberous sclerosis complex. *Pediatrics* 2006;118:1146-1151.
- Hancock E, Osborne J. Lymphangioliomyomatosis: a review of the literature. *Respir Med* 2002;96:1-6.
- O'Callaghan FJ, Noakes MJ, Martyn C, Osborne JP. An epidemiological study of renal pathology in tuberous sclerosis complex. *BJU Int* 2004;94:853-857.
- Brook-Carter P, Peral B, Ward CJ, Thompson P, Hughes J,

- Maheshwar MM, et al. Deletion of the TSC2 and pKD1 genes associated with severe infantile polycystic kidney disease—a contiguous gene syndrome. *Nat Genet* 1994;8:328-332.
- 26 Rose V, Au K, Pollom G, Roach ES, Prashner HR, Northrup H. Germ-line mosaicism in tuberous sclerosis: how common? *Am J Hum Genet* 1999;64:986-992.
- 27 Fryer A, Chalmers A, Connor J, Fraser I, Povey S, Yates AD, et al. Evidence that the gene for tuberous sclerosis is on chromosome 9. *Lancet* 1987;1:659-661.
- 28 Kandt R, Haines JI, Smith M, Northrup H, Gardner RJ, Short MP, et al. Linkage of an important gene locus for tuberous sclerosis to a chromosome 16 marker for polycystic kidney disease. *Nat Genet* 1992;2:37-41.
- 29 van Slegtenhorst M, de Hoogt R, Hermans C, Nellist M, Janssen B, Verhoef S, et al. Identification of the tuberous sclerosis gene *TSC1* on chromosome 9q34. *Science* 1997;277:805-808.
- 30 Dabora SL, Jozwiak S, Franz DN, Roberts PS, Nieto A, Chung J, et al. Mutational analysis in a cohort of 224 tuberous sclerosis patients indicates increased severity of TSC2, compared with TSC1, disease in multiple organs. *Am J Hum Genet* 2001;68:64-80.
- 31 Jones AC, Daniells CE, Snell RG, Tachataki M, Idziaszczyk SA, Krawczak M, et al. Molecular genetic and phenotypic analysis reveals differences between TSC1 and TSC2 associated familial and sporadic tuberous sclerosis. *Hum Mol Genet* 1997;6:2155-2161.
- 32 Kwiatkowski J, Wigowska-Sowinska J, Napierala D, Slomski R, Kwiatkowski DJ. Mosaicism in tuberous sclerosis as a potential cause of the failure of molecular diagnosis. *N Engl J Med* 1999;340:703-707.
- 33 van Slegtenhorst M, Nellist M, Nagelkerken B, Cheadle J, Snell R, van den Ouweland A, et al. Interaction between hamartin and tuberin, the TSC1 and TSC2 gene products. *Hum Mol Genet* 1998;7:1053-1057.
- 34 Kwiatkowski DJ. Rhebbing up mTor: new insights on TSC1 and TSC2, and the pathogenesis of tuberous sclerosis. *Cancer Biol Ther* 2003;2:471-476.
- 35 Franz DN, Leonard J, Tudor C, Chuck G, Care M, Sethuraman G, et al. Rapamycin causes regression of astrocytomas in tuberous sclerosis complex. *Ann Neurol* 2006;59:490-498.
- 36 Koenig MK, Butler IJ, Northrup H. Regression of subependymal giant cell astrocytoma with rapamycin in tuberous sclerosis complex. *J Child Neurol* 2008;23:1238-1239.
- 37 Sinson G, Sutton L, Yachnis A. Subependymal giant cell astrocytomas in children. *Pediatr Neurosurg* 1994;20:233-239.
- 38 Shepherd CW, Scheithauer BW, Gomez MR, Altermatt HJ, Katzmann JA. Subependymal giant cell tumor: a clinical, pathological, and flow cytometric study. *Neurosurgery* 1991;28:864-868.
- 39 Chow CW, Klug GL, Lewis EA. Subependymal giant-cell astrocytoma in children. An unusual discrepancy between histological and clinical features. *J Neurosurg* 1988;68:880-883.
- 40 Gyure KA, Prayson RA. Subependymal giant cell astrocytoma: a clinicopathologic study with HMB45 and MIB-1 immunohistochemical analysis. *Mod Pathol* 1997;10:313-317.
- 41 Knudson AG Jr. Mutation and cancer: statistical study of retinoblastoma. *Proc Natl Acad Sci USA* 1971;68:820-823.
- 42 Henske EP, Scheithauer BW, Short MP, Wollmann R, Nahmias J, Hornigold N, et al. Allelic loss is frequent in tuberous sclerosis kidney lesions but rare in brain lesions. *Am J Hum Genet* 1996;59:400-406.
- 43 Sepp T, Yates Jr, Green AJ. Loss of heterozygosity in tuberous sclerosis hamartomas. *J Med Genet* 1996;33:962-964.
- 44 Rakowski SK, Winterkorn EB, Paul E, Steele DJ, Halpern EF, Thiele EA. Renal manifestations of tuberous sclerosis complex: incidence, prognosis, and predictive factors. *Kidney Int* 2006;70:1777-1782.
- 45 Shepherd CW, Gomez MR, Lie JT, Crowson CS. Causes of death in patients with tuberous sclerosis. *Mayo Clin Proc* 1991;66:792-796.
- 46 Henske EP, Neumann HP, Scheithauer BW, Herbst EW, Short MP, Kwiatkowski DJ. Loss of heterozygosity in the tuberous sclerosis (*TSC2*) region of chromosome band 16p13 occurs in sporadic as well as TSC-associated renal angiomyolipomas. *Genes Chromosomes Cancer* 1995;13:295-298.
- 47 Han S, Santos TM, Puga A, Roy J, Thiele EA, McCollin M, et al. Phosphorylation of tuberin as a novel mechanism for somatic inactivation of the tuberous sclerosis complex proteins in brain lesions. *Cancer Res* 2004;64:812-816.
- 48 Carbonara C, Longa L, Grosso E, Borrone C, Garrè MG, Brisigotti M, et al. 9q34 loss of heterozygosity in a tuberous sclerosis astrocytoma suggests a growth suppressor-like activity also for the *TSC1* gene. *Hum Mol Genet* 1994;3:1829-1832.
- 49 Frerebeau PH, Benezech J, Segnarbieux F, Harbi H, Desy A, Marty-Double C. Intraventricular tumors in tuberous sclerosis. *Childs Nerv Syst* 1985;1:45-48.
- 50 Raju GP, Urion DK, Sahin M. Neonatal subependymal giant cell astrocytoma: new case and review of the literature. *Pediatr Neurol* 2007;36:128-131.
- 51 Scheithauer BW, Reagan TJ. Neuropathology. In: Gomez MR, Sampson JR, Whittemore VH, eds. *Tuberous sclerosis complex: developmental perspectives in psychiatry*, 3rd ed. New York: Oxford University Press, 1999: 101-144.
- 52 Nagib MG, Haines SJ, Erickson DL, Mastro AR. Tuberous sclerosis: a review for the neurosurgeon. *Neurosurgery* 1984;14:93-98.
- 53 Waga S, Yamamoto Y, Kojima T, Sakakura M. Massive haemorrhage in tumor of tuberous sclerosis. *Surg Neurol* 1977;8:99-101.
- 54 Fujiwara S, Takaki T, Hikita T, Nishio S. Subependymal giant-cell astrocytoma associated with tuberous sclerosis. Do subependymal nodules grow? *Childs Nerv Syst* 1989;5:43-44.
- 55 Holanda FJ, Holanda GM. Tuberous sclerosis—neurosurgical indications in intraventricular tumors. *Neurosurg Rev* 1980;3:139-150.
- 56 Torres OA, Roach ES, Delgado MR, Sparagana SP, Sheffield E, Swift D, et al. Early diagnosis of subependymal giant cell astrocytoma in patients with tuberous sclerosis. *J Child Neurol* 1998;13:173-177.
- 57 Kumar R, Singh V. Subependymal giant cell astrocytomas: a report of five cases. *Neurosurg Rev* 2004;27:274-280.
- 58 Sharma MC, Ralte AM, Gaekwad S, Santosh V, Shankar SK, Sarkar C. Subependymal giant cell astrocytoma—a clinicopathological study of 23 cases with special emphasis on histogenesis. *Pathol Oncol Res* 2004;10:219-224.
- 59 Osborne JP, Merrifield J, O'Callaghan FJ. Tuberous sclerosis—what's new? *Arch Dis Child* 2008;93:728-731.
- 60 Franz DN, de Vries PJ, Crino PB. Giant cell astrocytomas in tuberous sclerosis complex. *Arch Dis Child* 2008;93:751-754.

Received May 12, 2009

Accepted after revision February 9, 2010