

Identifying key genes associated with Hirschsprung's disease based on bioinformatics analysis of RNA-sequencing data

Wei-Kang Pan, Ya-Fei Zhang, Hui Yu, Ya Gao, Bai-Jun Zheng, Peng Li, Chong Xie, Xin Ge
Xi'an, China

Background: Hirschsprung's disease (HSCR) is a type of megacolon induced by deficiency or dysfunction of ganglion cells in the distal intestine and is associated with developmental disorders of the enteric nervous system. To explore the mechanisms of HSCR, we analyzed the RNA-sequencing data of the expansion and the narrow segments of colon tissues separated from children with HSCR.

Methods: RNA-sequencing of the expansion segments and the narrow segments of colon tissues isolated from children with HSCR was performed. After differentially expressed genes (DEGs) were identified using the edgeR package in R, functional and pathway enrichment analyses of DEGs were carried out using DAVID software. To further screen the key genes, protein-protein interaction (PPI) network and module analyses were conducted separately using Cytoscape software.

Results: A total of 117 DEGs were identified in the expansion segment samples, including 47 up-regulated and 70 down-regulated genes. Functional enrichment analysis suggested that *FOS* and *DUSP1* were implicated in response to endogenous stimulus. In the PPI network analysis, *FOS* (degree=20), *EGR1* (degree=16), *ATF3* (degree=9), *NOS1* (degree=8), *CCL5* (degree=8), *DUSP1* (degree=7), *CXCL3* (degree=6), *VIP* (degree=6), *FOSB* (degree=5), and *NOS2* (degree=4) had higher degrees, which could interact with other genes. In addition, two significant modules (module 1 and module 2) were identified from the PPI network.

Conclusion: Several genes (including *FOS*, *EGR1*, *ATF3*, *NOS1*, *CCL5*, *DUSP1*, *CXCL3*, *VIP*, *FOSB*, and

NOS2) might be involved in the development of HSCR through their effect on the nervous system.

World J Pediatr 2017;13(3):267-273

Key words: differentially expressed genes; functional and pathway enrichment analysis; Hirschsprung's disease; protein-protein interaction network

Introduction

Hirschsprung's disease (HSCR), also called Hirschsprung disease, congenital aganglionic megacolon or congenital megacolon, is a type of megacolon induced by deficiency or dysfunction of ganglion cells in the distal intestine.^[1] In HSCR, the affected colon segment cannot relax the bowels; therefore, HSCR is characterized by fecal retention, intestinal obstruction, constipation and abdominal distention.^[2] HSCR usually occurs in children, especially boys, and seriously impacts quality of life and, possibly, mortality.^[3] In Japan, the incidence rate of HSCR is approximately one in 5000 births.^[4] Treatment strategies of HSCR include resection of the abnormal colon segment and reanastomosis. However, some children still have bowel dysfunction after treatment and need a lifetime of intestinal fistulas.^[5] Thus, it is necessary to investigate the mechanisms of HSCR and to develop novel targeted treatment strategies for this disease.

To date, several studies have explored the key genes involved in HSCR. For instance, the *RET* gene, which encodes a tyrosine-kinase receptor, plays an important role in the development of HSCR by affecting the enteric ganglia.^[6] Previous studies have demonstrated that neuregulin 1 (*NRG1*) can serve as an HSCR susceptibility locus via mediating the development of the enteric ganglia precursors.^[7,8] The ablation of glial cell line-derived neurotrophic factor (*GDNF*) family receptor α_1 (*GFR α_1*), which is a high affinity receptor for *GDNF*, can cause rapid and extensive death of enteric neurons and result in colon aganglionosis reminiscent of HSCR in mice during late gestation.^[9] Neurotrophin

Author Affiliations: Department of Pediatric Surgery, The Second Affiliated Hospital of Xi'an Jiaotong University, Xi'an 710004, China (Pan WK, Yu H, Gao Y, Zheng BJ, Li P, Xie C, Ge X); Department of Endoscopy, Shaanxi Nuclear Industry 215 Hospital, Xianyang 712000, China (Zhang YF)

Corresponding Author: Ya Gao, Department of Pediatric Surgery, The Second Affiliated Hospital of Xi'an Jiaotong University, Xi'an 710004, China (Tel: +86-2987679373; Fax: +86-2987679373; Email: ygao@mail.xjtu.edu.cn)

doi: 10.1007/s12519-017-0002-0

Online First, January 2017

©Children's Hospital, Zhejiang University School of Medicine, China and Springer-Verlag Berlin Heidelberg 2017. All rights reserved.

3 (NTF3) and neurotrophic tyrosine kinase receptor 3 (NTRK3) may function in the pathogenesis of HSCR by affecting the development of the enteric nervous system (ENS) through the NTF3/NTRK3 signaling pathway.^[10,11] Duan et al^[12] found mutations of the endothelin 3 (*EDN-3*) and endothelin receptor type B (*EDNRB*) genes in short-segment HSCR in the Chinese population and determined that these genes may be correlated with the pathogenesis of HSCR. Cantrell et al^[13] reported that interaction between sex determining region Y (SRY)-box 10 (Sox10) and endothelin receptor type B (EdnrB) affects the development of the ENS in mouse models and can affect the epistatic network, producing phenotypic variation among patients with aganglionosis.

RNA-sequencing (or massively parallel cDNA sequencing), which can detect gene changes, gene fusion, and somatic mutation, is an accurate method for quantifying and characterizing transcriptomes.^[14] Using RNA-sequencing data, Saeed et al^[15] analyzed differentially expressed genes (DEGs) between abnormal and normal bowel segments in patients with HSCR relative to controls and found that reelin (RELN), galanin/GMAP prepropeptide (GAL), growth associated protein 43 (GAP43), Neurexin-1 (NRSN1), and gamma 2 subunit of the gamma-aminobutyric acid receptor type A (GABRG2) correlate with the pathogenesis of HSCR. It is proposed that different analysis processes and thresholds can generate different results.^[16] To gain a more comprehensive understanding of the pathogenesis of HSCR, RNA-sequencing data were used to analyze DEGs between expansion and narrow segment samples. In addition, enrichment analysis, protein-protein interaction (PPI) network construction and module analysis were carried out separately to screen critical genes in HSCR.

Methods

Sample collection

This study was approved by the Institutional Review Board of the Second Affiliated Hospital of Xi'an Jiaotong University. In addition, all participants provided their informed consent. Using electronic colonoscopy, expansion segments and narrow segments of colon tissues were excised from children with HSCR who were treated in the Second Affiliated Hospital of Xi'an Jiaotong University from September 2014 to July 2015. The expansion segments and the narrow segments were randomly mixed in their respective groups to form 2 sequencing samples. All the patients were diagnosed with HSCR by barium enema and anorectal manometry evaluation before surgical treatment. Before surgery, a suction rectal biopsy was obtained to verify

aganglionosis. After surgery, pathological staining was performed to establish a definitive diagnosis. The isolated tissues were cut into small pieces (<0.5 cm) and soaked in 5 times the volume of RNAlater (Ambion, Foster City, CA, USA) at 4°C overnight then stored at -80°C.

RNA extraction and RNA-sequencing library construction

Total RNA was extracted from the expansion and the narrow segments of colon tissues using TRIzol reagent (Invitrogen, Shanghai, China) according to the manufacturer's instructions. The integrity and purity of total RNA were detected using 2% agarose gel electrophoresis and spectrophotometry (Merinton, Beijing, China), respectively. Finally, RNA-seq library construction and high-throughput sequencing were separately carried out using a NEBNext[®] Ultra[™] RNA Library Prep Kit for Illumina[®] (New England Biolabs, Ipswich, MA, USA) and an Illumina HiSeq 2500 v4 100PE System (Illumina, San Diego, CA, USA).

Data preprocessing and DEG screening

The NGSQC Toolkit^[17] was used to preprocess the paired-end sequencing data and filter out low quality sequences. Using TopHat2 software,^[18] sequence alignment of the preprocessed data was performed with the hg19 human genome sequences from the UCSC website (<http://genome.ucsc.edu>) as references.^[19] Subsequently, the expression levels of genes were calculated using Cuffdiffs software^[20] to obtain a gene expression matrix of samples. The "edgeR" package^[21] in R was used to screen for DEGs between the expansion and the narrow segment samples. The thresholds were P value <0.05 and $|\log_2 \text{fold change (FC)}| > 1$.

Functional and pathway enrichment analyses

Gene Ontology (GO, <http://www.geneontology.org/>) seeks to develop and use a series of controlled, structured vocabularies for annotating sequences, genes and their products.^[22] The Kyoto Encyclopedia of Genes and Genomes (KEGG, <http://www.genome.jp/kegg/>) aims to integrate biological systems information of genomic, chemical and systemic functional aspects.^[23] Using DAVID software, GO functional and KEGG pathway enrichment analyses of DEGs were performed. Terms involving more than 2 genes and with a P value <0.05 were selected.

PPI network construction and module analysis

The Search Tool for the Retrieval of Interacting Genes (STRING, <http://string-db.org/>) database gives a critical evaluation and integration of PPIs consisting of functional and physical associations.^[24] Using the STRING database,^[24] the PPI relationships among the DEGs were analyzed, with the required confidence (combined score) >0.4 as the threshold. After the PPIs were searched, the PPI network

was constructed using Cytoscape software (<http://www.cytoscape.org/>).^[25] In the PPI network, nodes represented proteins, and their degrees were equal to the number of edges linked to them. Moreover, module analysis was performed for the PPI network using the MCODE plugin^[26] in Cytoscape with the default parameters.

Results

Analysis of DEGs

DEGs with a *P* value <0.05 and $|\log_2FC| >1$ between the expansion and the narrow segment samples were

analyzed. Compared with narrow segment samples, 117 DEGs were identified in the expansion segment samples, including 47 up-regulated and 70 down-regulated genes.

Functional and pathway enrichment analyses

Using the DAVID software, enrichment analyses were carried out on the up-regulated genes and the down-regulated genes separately. The up-regulated genes were significantly enriched in 43 GO terms, and the top 10 GO terms are listed in Table 1. The top 10 GO terms included response to endogenous stimulus (*P*<0.001), which involved FBJ osteosarcoma oncogene (*FOS*)

Table 1. The top 10 Gene Ontology (GO) functions enriched for the up-regulated genes

Description	<i>P</i> value	Gene number	Gene symbol
GO:0009719-response to endogenous stimulus	<0.001	9	<i>ME1, FOS, DUSP1, SOCS3, HMGCS1, NR4A2, GAL, JUNB, MGST1</i>
GO:0010033-response to organic substance	<0.001	11	<i>ME1, EGR1, FOS, DUSP1, SOCS3, HMGCS1, NR4A2, GAL, JUNB, CHRNA3, MGST1</i>
GO:0042493-response to drug	<0.001	7	<i>FOS, SOCS3, HMGCS1, GAL, JUNB, CHRNA3, MGST1</i>
GO:0032570-response to progesterone stimulus	<0.001	3	<i>FOS, SOCS3, JUNB</i>
GO:0007610-behavior	<0.001	7	<i>EGR1, FOS, CXCL3, NR4A2, FOSB, GAL, CHRNA3</i>
GO:0048545-response to steroid hormone stimulus	<0.001	5	<i>FOS, DUSP1, SOCS3, GAL, JUNB</i>
GO:0009314-response to radiation	<0.001	5	<i>FOS, BLM, DUSP1, SOCS3, JUNB</i>
GO:0007167-enzyme linked receptor protein signaling pathway	<0.001	6	<i>FOS, HBEGF, AREG, GDF15, CHRNA3, TOB1</i>
GO:0009725-response to hormone stimulus	<0.001	6	<i>ME1, FOS, DUSP1, SOCS3, GAL, JUNB</i>
GO:0009628-response to abiotic stimulus	<0.001	6	<i>ME1, FOS, DUSP1, SOCS3, HMGCS1, NR4A2, GAL, JUNB, MGST1</i>

Table 2. The top 10 Gene Ontology (GO) functions and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways enriched for the down-regulated genes

Description	<i>P</i> value	Gene number	Gene symbol
The top 10 GO functions enriched for the down-regulated genes			
GO:0006959-humoral immune response	<0.001	8	<i>CIQA, CIQB, C7, C3, FCN3, PAX5, CFD, EB13</i>
GO:0002526-acute inflammatory response	<0.001	8	<i>CIQA, CIQB, C7, C3, FCN3, SERPINA3, VNN1, CFD</i>
GO:0006955-immune response	<0.001	15	<i>C7, HLA-DRB1, C3, TNFRSF13B, TNFRSF17, PAX5, CCL5, CIQA, CIQB, FCN3, VNN1, NOS2, CFD, EB13, CD27</i>
GO:0002684-positive regulation of immune system process	<0.001	10	<i>CIQA, CIQB, C7, C3, FCN3, VNN1, NOS2, CFD, CD27, EB13</i>
GO:0006956-complement activation	<0.001	6	<i>CIQA, CIQB, C7, C3, FCN3, CFD</i>
GO:0002541-activation of plasma proteins involved in acute inflammatory response	<0.001	6	<i>CIQA, CIQB, C7, C3, FCN3, CFD</i>
GO:0045087-innate immune response	<0.001	8	<i>CIQA, CIQB, C7, C3, FCN3, VNN1, NOS2, CFD</i>
GO:0006954-inflammatory response	<0.001	10	<i>CIQA, KNG1, CIQB, C7, C3, FCN3, SERPINA3, VNN1, CFD, CCL5</i>
GO:0002250-adaptive immune response	<0.001	6	<i>CIQA, CIQB, C7, C3, CD27, EB13</i>
GO:0002460-adaptive immune response based on somatic recombination of immune receptors built from immunoglobulin superfamily domains	<0.001	6	<i>CIQA, CIQB, C7, C3, CD27, EB13</i>
The KEGG pathways enriched for the down-regulated genes			
hsa04610-complement and coagulation cascades	<0.001	6	<i>CIQA, KNG1, CIQB, C7, C3, CFD</i>
hsa05020-prion diseases	<0.001	4	<i>CIQA, CIQB, C7, CCL5</i>
hsa05322-systemic lupus erythematosus	<0.05	5	<i>CIQA, CIQB, C7, HLA-DRB1, C3</i>
hsa04672-intestinal immune network for IgA production	<0.05	4	<i>HLA-DRB1, TNFRSF13B, CCR10, TNFRSF17</i>
hsa00980-metabolism of xenobiotics by cytochrome P450	<0.05	3	<i>GSTM1, ADH1B, GSTM5</i>
hsa00982-drug metabolism	<0.05	3	<i>GSTM1, ADH1B, GSTM5</i>
hsa04060-cytokine-cytokine receptor interaction	<0.05	5	<i>TNFRSF13B, CCR10, TNFRSF17, CCL5, CD27</i>

and dual specificity phosphatase 1 (*DUSP1*), response to organic substance ($P<0.001$) and response to drug ($P<0.001$). No pathways were enriched among the up-regulated genes.

The down-regulated genes were significantly enriched in 51 GO terms, including humoral immune

response ($P<0.001$), acute inflammatory response ($P<0.001$) and immune response ($P<0.001$) (Table 2). Meanwhile, 7 pathways were enriched among the down-regulated genes, including complement and coagulation cascades ($P<0.001$), prion diseases ($P<0.001$), which involved CC chemokine ligand 5




Fig. 1. The protein-protein interaction (PPI) network of the differentially expressed genes (DEGs). Red and green nodes represent up-regulated genes and down-regulated genes, respectively. Different shades of nodes indicate the degree of significance of differential expression.

Table 3. The top 10 Gene Ontology (GO) functions enriched for the genes involved in module 1 and module 2

Module	Description	P value	Gene number	Gene symbol
Module 1	GO:0007186-G-protein coupled receptor protein signaling pathway	<0.001	8	<i>VIP, KNG1, C3, CXCL3, CCR10, GAL, CCL5, SST</i>
	GO:0007166-cell surface receptor linked signal transduction	<0.001	8	<i>VIP, KNG1, C3, CXCL3, CCR10, GAL, CCL5, SST</i>
	GO:0006954-inflammatory response	<0.001	5	<i>KNG1, C3, CXCL3, GAL, CCL5</i>
	GO:0009611-response to wounding	<0.001	5	<i>KNG1, C3, CXCL3, GAL, CCL5</i>
	GO:0006952-defense response	<0.001	5	<i>KNG1, C3, CXCL3, GAL, CCL5</i>
	GO:0046903-secretion	<0.001	4	<i>VIP, KNG1, GAL, CCL5</i>
	GO:0007610-behavior	<0.05	4	<i>CXCL3, CCR10, GAL, CCL5</i>
	GO:0032101-regulation of response to external stimulus	<0.05	3	<i>KNG1, C3, CCL5</i>
Module 2	GO:0006935-chemotaxis	<0.05	3	<i>CXCL3, CCR10, CCL5</i>
	GO:0006357-regulation of transcription from RNA polymerase II promoter	<0.001	5	<i>EGR1, NR4A2, NR4A1, FOSB, JUNB</i>
	GO:0006355-regulation of transcription, DNA-dependent	<0.001	6	<i>EGR1, ATF3, NR4A2, NR4A1, FOSB, JUNB</i>
	GO:0051252-regulation of RNA metabolic process	<0.001	6	<i>EGR1, ATF3, NR4A2, NR4A1, FOSB, JUNB</i>
	GO:0045944-positive regulation of transcription from RNA polymerase II promoter	<0.001	4	<i>EGR1, NR4A2, NR4A1, JUNB</i>
	GO:0045893-positive regulation of transcription, DNA-dependent	<0.001	4	<i>EGR1, NR4A2, NR4A1, JUNB</i>
	GO:0051254-positive regulation of RNA metabolic process	<0.001	4	<i>EGR1, NR4A2, NR4A1, JUNB</i>
	GO:0045941-positive regulation of transcription	<0.05	4	<i>EGR1, NR4A2, NR4A1, JUNB</i>
	GO:0045449-regulation of transcription	<0.05	6	<i>EGR1, ATF3, NR4A2, NR4A1, FOSB, JUNB</i>
	GO:0010628-positive regulation of gene expression	<0.05	4	<i>EGR1, NR4A2, NR4A1, JUNB</i>
GO:0045935-positive regulation of nucleobase, nucleoside, nucleotide and nucleic acid metabolic process	<0.05	4	<i>EGR1, NR4A2, NR4A1, JUNB</i>	




Fig. 2. The two significant modules (module 1 and module 2) identified from the protein-protein interaction (PPI) network. Red and green nodes represent up-regulated genes and down-regulated genes, respectively. Different shades of nodes indicate the degree of significance of differential expression.

(CCL5), and systemic lupus erythematosus ($P<0.005$) (Table 2).

PPI network construction and module analysis

The PPI network of the DEGs had 59 nodes and 113 interactions (Fig. 1). In particular, FOS (degree=20), early growth response 1 (EGR1, degree=16), activating transcription factor 3 (ATF3, degree=9), nitric oxide synthase 1 (NOS1, degree=8), CCL5 (degree=8), DUSP1 (degree=7), CXCL3 (degree=6), vasoactive intestinal peptide (VIP, degree=6), FBJ murine osteosarcoma viral oncogene homolog B (FOSB, degree=5), and nitric oxide synthase 2 (NOS2, degree=4) had higher degrees and could interact with others genes in the PPI network. Afterwards, module analysis was performed for the PPI network, and two significant modules (module 1 and module 2) were identified (Fig. 2). The top 10 enriched functions among the genes in module 1 are shown in Table 3 and include G-protein coupled receptor protein signaling pathway ($P<0.001$), cell surface receptor linked signal transduction ($P<0.001$) and taxis ($P<0.005$), which involved CXCL3 and CCL5. The genes in module 2 were enriched in functions such as regulation of transcription from RNA polymerase II promoter ($P<0.001$), DNA-dependent regulation of transcription ($P<0.001$) and regulation of RNA metabolic process ($P<0.001$) (Table 3).

Discussion

As the main component of the peripheral nervous system, the ENS is a complex network of glia and neurons within the intestinal wall,^[27,28] and a developmental disorder of the ENS has been correlated with HSCR.^[29] In this study, 117 DEGs were identified in the expansion segment samples, including 47 up-regulated and 70 down-regulated genes. In the PPI network, FOS (degree=20), EGR1 (degree=16), ATF3 (degree=9), NOS1 (degree=8), CCL5 (degree=8), DUSP1 (degree=7),

CXCL3 (degree=6), VIP (degree=6), FOSB (degree=5), and NOS2 (degree=4) had higher degrees, which had interactions with other genes. Moreover, two significant modules (module 1 and module 2) were identified in the PPI network.

Functional enrichment analysis showed that *FOS* and *DUSP1* were implicated in response to endogenous stimulus. The *c-fos* gene is related to apoptosis and regeneration of damaged retinal ganglion cells (RGCs), indicating that the *c-fos*-mediated cascade precisely correlates with apoptosis and that regeneration of RGC plays a critical role in neuronal survival and regeneration.^[30] As a member of the Fos family, FosB is up-regulated after an acute increase in central venous pressure in unanesthetized rats.^[31] DUSP1 can specially regulate sensory input neurons of the telencephalon and thalamus, which may reduce stimulus-induced expression of immediate early genes such as *EGR1*.^[32] These findings suggest that *FOS*, *FOSB*, *DUSP1* and *EGR1* might be correlated with the development of HSCR.

Enrichment analyses indicated that CCL5 was involved in the pathway of prion diseases, while CXCL3 and CCL5 were functionally enriched in taxis. Inhibition of CCL5 by neutralizing mAb decreased the severity of central nervous system (CNS) disease in a viral model of demyelination.^[33] CC receptor 5 (CCR5) and its primary ligand CCL5 play critical roles in promoting the transport of leukocytes into the CNS of mice infected by mouse hepatitis virus (MHV).^[34-36] CXCL3 has a low expression level in 12-O-tetradecanoyl phorbol-13-acetate inducible sequence 21 (Tis21)-null cerebellar granule neuron precursor cells (GCPs) of lesions and the external granular layer, and CXCL3 activation induced by Tis21 contributes to migration of GCPs.^[37] Therefore, CCL5 and CXCL3 might be implicated in the pathogenesis of HSCR.

In addition, as a member of the ATF/cyclic AMP responsive element binding family of transcription factors, ATF3 is activated in motoneurons and sensory neurons in the spinal cord after nerve injury and may

be a promising marker of nerve injury in the nervous system.^[38] ATF3 can promote neurite elongation of dorsal root ganglion, indicating that ATF3 helps nerve regeneration by improving the intrinsic growth state of injured neurons.^[39-42] VIP is a neurotransmitter in the mucosa, lamina propria, and myenteric plexus of the small and large intestine, which can respond to secretomotor reflexes and activate adenylate cyclase.^[43] A previous study has shown that VIP functions in reducing the net intestinal fluid absorption in patients with chemical colitis, which can be completely inhibited by its antagonist [4Cl-d-Phe6, Leu17]VIP.^[44] Nitric oxide synthases (NOS) are the predominant enzymes producing nitric oxide (NO) from conversion of L-arginine and generally need NADPH and oxygen co-substrates.^[45] Additionally, NO produced by the enteric nervous system is conducive for maintaining gut homeostasis through regulation of vascular tone, gut motility, mucosal barrier function, blood supply, immunity and secretions.^[46] These findings suggest that *ATF3*, *VIP*, *NOS1* and *NOS2* might also function in HSCR by affecting the nervous system.

In conclusion, based on the bioinformatics analyses, 117 DEGs were identified in the expansion segment samples. Moreover, several genes (including *FOS*, *EGRI*, *ATF3*, *NOS1*, *CCL5*, *DUSP1*, *CXCL3*, *VIP*, *FOSB*, and *NOS2*) might be involved in the development of HSCR through their impact on the nervous system. However, these predicted results must be further confirmed through experimental studies, such as quantitative real-time polymerase chain reaction (qRT-PCR) in the future.

Funding: This work was supported by the National Natural Science Foundation of China (NSFC) (No. 30872699), and Natural Science Basic Research Plan in Shaanxi Province of China (No. 2016JQ8050).

Ethical approval: The study protocol was approved by the Scientific Research Ethics Committee at Xi'an Jiaotong University Health Science Center (no. 2008032) in accordance with the latest version of the Declaration of Helsinki and the national legal regulations.

Competing interest: There is no competing interest.

Contributors: PWK and ZYF carried out the conception and design of the research; YH, GY participated in the acquisition of data; ZBJ, LP carried out the analysis and interpretation of data; XC and GX participated in the design of the study and performed the statistical analysis. All authors read and approved the final manuscript.

References

- Puri P, Shinkai T. Pathogenesis of Hirschsprung's disease and its variants: recent progress. *Semin Pediatr Surg* 2004;13:18-24.
- De LF, Boeckxstaens GE, Benninga MA. Symptomatology, pathophysiology, diagnostic work-up, and treatment of Hirschsprung disease in infancy and childhood. *Curr Gastroenterol Rep* 2007;9:245-253.
- Butler Tjaden NE, Trainor PA. The developmental etiology and pathogenesis of Hirschsprung disease. *Transl Res* 2013;162:1-15.
- Suita S, Taguchi T, Ieiri S, Nakatsuji T. Hirschsprung's disease in Japan: analysis of 3852 patients based on a nationwide survey in 30 years. *J Pediatr Surg* 2005;40:197-202.
- Okamura Y, Saga Y. Notch signaling is required for the maintenance of enteric neural crest progenitors. *Development* 2008;135:3555-3565.
- de Pontual L, Pelet A, Trochet D, Jaubert F, Espinosa-Parrilla Y, Munnich A, et al. Mutations of the RET gene in isolated and syndromic Hirschsprung's disease in human disclose major and modifier alleles at a single locus. *J Med Genet* 2006;43:419-423.
- Garcia-Barcelo MM, Tang CS, Ngan ES, Lui VC, Chen Y, So MT, et al. Genome-wide association study identifies NRG1 as a susceptibility locus for Hirschsprung's disease. *Proc Natl Acad Sci U S A* 2009;106:2694-2699.
- Phusantisampan T, Sangkhathat S, Phongdara A, Chiengkriwate P, Patrapinyokul S, Mahasirimongkol S. Association of genetic polymorphisms in the RET-protocogene and NRG1 with Hirschsprung disease in Thai patients. *J Hum Genet* 2012;57:286-293.
- Uesaka T, Jain S, Yonemura S, Uchiyama Y, Milbrandt J, Enomoto H. Conditional ablation of GFR α 1 in postmigratory enteric neurons triggers unconventional neuronal death in the colon and causes a Hirschsprung's disease phenotype. *Development* 2007;134:2171-2181.
- Fernández RM, Sánchez-Mejías A, Mena M, Ruiz-Ferrer M, López-Alonso M, Antiñolo G, et al. A novel point variant in NTRK3, R645C, suggests a role of this gene in the pathogenesis of Hirschsprung disease. *Ann Hum Genet* 2009;73:19-25.
- Ruiz-Ferrer M, Fernandez RM, Antiñolo G, Lopez-Alonso M, Borrego S. NTF-3, a gene involved in the enteric nervous system development, as a candidate gene for Hirschsprung disease. *J Pediatr Surg* 2008;43:1308-1311.
- Duan X-L, Zhang X-S, Li G-W. Clinical relationship between EDN-3 gene, EDNRB gene and Hirschsprung's disease. *World J Gastroenterol* 2003;9:2839.
- Cantrell VA, Owens SE, Chandler RL, Airey DC, Bradley KM, Smith JR, et al. Interactions between Sox10 and EdnrB modulate penetrance and severity of aganglionosis in the Sox10Dom mouse model of Hirschsprung disease. *Hum Mol Genet* 2004;13:2289-2301.
- Ozsolak F, Milos PM. RNA sequencing: advances, challenges and opportunities. *Nat Rev Genet* 2011;12:87-98.
- Saeed A, Barreto L, Neogii SG, Loos A, Mcfarlane I, Aslam A. Identification of novel genes in Hirschsprung disease pathway using whole genome expression study. *J Pediatr Surg* 2012;47:303-307.
- Afsari B, Geman D, Fertig EJ. Learning dysregulated pathways in cancers from differential variability analysis. *Cancer Inform* 2014:61.
- Patel RK, Jain M. NGS QC Toolkit: a toolkit for quality control of next generation sequencing data. *PLoS One* 2012;7:e30619.
- Kim D, Perteau G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome Biol* 2013;14:R36.
- Karolchik D, Baertsch R, Diekhans M, Furey TS, Hinrichs A, Lu Y, et al. The UCSC genome browser database. *Nucleic Acids Res* 2003;31:51-54.

- 20 Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, et al. Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nat Protoc* 2012;7:562-578.
- 21 Smyth GK. Limma: linear models for microarray data. In: Gentleman R, Carey VJ, Huber W, Irizarry RA, Dudoit S, eds. *Bioinformatics and computational biology solutions using R and Bioconductor*. New York: Springer, 2005: 397-420.
- 22 Consortium GO. The Gene Ontology in 2010: extensions and refinements. *Nucleic acids research* 2010;38:D331-D335.
- 23 Kanehisa M, Araki M, Goto S, Hattori M, Hirakawa M, Itoh M, et al. KEGG for linking genomes to life and the environment. *Nucleic Acids Res* 2008;36:D480-D484.
- 24 Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, et al. STRING v10: protein-protein interaction networks, integrated over the tree of life. *Nucleic Acids Res* 2015;43:D447-D52.
- 25 Saito R, Smoot ME, Ono K, Ruscheinski J, Wang P-L, Lotia S, et al. A travel guide to Cytoscape plugins. *Nat Methods* 2012;9:1069-1076.
- 26 Bader GD, Hogue CW. An automated method for finding molecular complexes in large protein interaction networks. *BMC Bioinformatics* 2003;4:2.
- 27 Furness JB. The enteric nervous system: normal functions and enteric neuropathies. *Neurogastroenterol Motil* 2008;20 Suppl 1:32-38.
- 28 Gershon MD. Developmental determinants of the independence and complexity of the enteric nervous system. *Trends Neurosci* 2010;33:446-456.
- 29 McKeown SJ, Stamp L, Hao MM, Young HM. Hirschsprung disease: a developmental disorder of the enteric nervous system. *Wiley Interdiscip Rev Dev Biol* 2013;2:113-129.
- 30 Oshitari T, Dezawa M, Okada S, Takano M, Negishi H, Horie H, et al. The role of c-fos in cell death and regeneration of retinal ganglion cells. *Invest Ophthalmol Vis Sci* 2002;43:2442-2449.
- 31 Howe BM, Bruno SB, Higgs KA, Stigers RL, Cunningham JT. FosB expression in the central nervous system following isotonic volume expansion in unanesthetized rats. *Exp Neurol* 2004;187:190-198.
- 32 Horita H, Wada K, Rivas MV, Hara E, Jarvis ED. The *dusp1* immediate early gene is regulated by natural stimuli predominantly in sensory input neurons. *J Comp Neurol* 2010; 518:2873-2901.
- 33 Glass WG, Hickey MJ, Hardison JL, Liu MT, Manning JE, Lane TE. Antibody targeting of the CC chemokine ligand 5 results in diminished leukocyte infiltration into the central nervous system and reduced neurologic disease in a viral model of multiple sclerosis. *J Immunol* 2004;172:4018-4025.
- 34 Glass WG, Liu MT, Kuziel WA, Lane TE. Reduced macrophage infiltration and demyelination in mice lacking the chemokine receptor CCR5 following infection with a neurotropic coronavirus. *Virology* 2001;288:8-17.
- 35 Glass WG, Lane TE. Functional expression of chemokine receptor CCR5 on CD4+ T cells during virus-induced central nervous system disease. *J Virol* 2003;77:191-198.
- 36 Glass WG, Lane TE. Functional analysis of the CC chemokine receptor 5 (CCR5) on virus-specific CD8+ T cells following coronavirus infection of the central nervous system. *Virology* 2003;312:407-414.
- 37 Farioli-Vecchioli S, Cinà I, Ceccarelli M, Micheli L, Leonardi L, Ciotti MT, et al. Tis 21 knock-out enhances the frequency of medulloblastoma in *Patched1* heterozygous mice by inhibiting the Cxcl3-dependent migration of cerebellar neurons. *J Neurosci* 2012;32:15547-15564.
- 38 Tsujino H, Kondo E, Fukuoka T, Dai Y, Tokunaga A, Miki K, et al. Activating transcription factor 3 (ATF3) induction by axotomy in sensory and motoneurons: A novel neuronal marker of nerve injury. *Mol Cell Neurosci* 2000;15:170-182.
- 39 Seiffers R, Mills CD, Woolf CJ. ATF3 increases the intrinsic growth state of DRG neurons to enhance peripheral nerve regeneration. *J Neurosci* 2007;27:7911-7920.
- 40 Hunt D, Raivich G, Anderson PN. Activating transcription factor 3 and the nervous system. *Front Mol Neurosci* 2012;5:7.
- 41 Seiffers R, Allchorne AJ, Woolf CJ. The transcription factor ATF-3 promotes neurite outgrowth. *Mol Cell Neurosci* 2006;32:143-154.
- 42 Pearson AG, Gray CW, Pearson JF, Greenwood JM, During MJ, Dragunow M. ATF3 enhances c-Jun-mediated neurite sprouting. *Mol Brain Res* 2003;120:38-45.
- 43 Waldman D, Gardner J, Zfass A, Makhlof G. Effects of vasoactive intestinal peptide, secretin, and related peptides on rat colonic transport and adenylate cyclase activity. *Gastroenterology* 1977;73:518-523.
- 44 Mourad FH, Barada KA, Rached NAB, Khoury CI, Saadé NE, Nassar CF. Inhibitory effect of experimental colitis on fluid absorption in rat jejunum: role of the enteric nervous system, VIP, and nitric oxide. *Am J Physiol-Gastrl* 2006;290:G262-G268.
- 45 Andrew PJ, Mayer B. Enzymatic function of nitric oxide synthases. *Cardiovasc Res* 1999;43:521-531.
- 46 Savidge TC. Importance of NO and its related compounds in enteric nervous system regulation of gut homeostasis and disease susceptibility. *Curr Opin Pharmacol* 2014;19:54-60.

Received March 23, 2016

Accepted after revision August 25, 2016